

Detection of Infectious Tobamoviruses in Pepper Seed

SEPTEMBER 2019

Developed by ISHI-Veg
All rights reserved - ©2019 ISF

Detection of Infectious Tobamoviruses in Pepper Seed

Crop: Pepper (*Capsicum annuum*)

Pathogen(s): *Tobacco mosaic virus* (TMV), *Tomato mosaic virus* (ToMV), *Tomato mottle mosaic virus* (ToMMV), *Tomato brown rugose fruit virus* (ToBRFV) and *Pepper mild mottle virus* (PMMoV; sometimes referred to as PMMV)

Note: For many years *Tobacco mosaic virus* (TMV), *Tomato mosaic virus* (ToMV) and *Pepper mild mottle virus* (PMMoV) have been the most prominently perceived tobamoviruses infecting pepper seed. Recently two other tobamoviruses related to but distinct from TMV and ToMV were characterized: *Tomato mottle mosaic virus* (ToMMV) (Li et al., 2013) and *Tomato brown rugose fruit virus* (ToBRFV) (Salem et al., 2016).

These viruses are shown to cause disease in pepper. As in the case of infection with TMV, ToMV and PMMoV, ToMMV (Sui et al., 2017) and ToBRFV (Luria et al., 2017) generate local lesions on leaves of indicator plants containing the N-gene (*Nicotiana tabacum* cv. Xanthi NN and *Nicotiana glutinosa*). Therefore, the method covers these two viruses as well.

Revision history: Version 6.3, September 2019

PRINCIPLE

Detection of infectious tobamoviruses in pepper seed by inoculating leaves of indicator plants *Nicotiana tabacum* cv. Xanthi NN or *Nicotiana glutinosa* with pepper seed extract. Infectious virions cause typical local lesions that demonstrate viability of the virus.

A DAS-ELISA assay may be used to pre-screen a seed extract. The test is complete if no virus is detected and the seed lot considered healthy. As ELISA detects both infectious virions and non-infectious virus particles, a positive ELISA is followed by a bioassay to determine the presence of infectious tobamovirus.




Figure 1. Method process flow

METHOD VALIDATION

The method was evaluated in a comparative test organized by ISHI-Veg. The US National Seed Health System (NSHS) (<http://seedhealth.org>) has approved the method as a Standard B.

RESTRICTIONS ON USE

This method is suitable for testing untreated seed.

It is also suitable for testing seed that has been treated using physical or chemical (acid extraction, calcium or sodium hypochlorite, tri-sodium phosphate, etc.) processes with the aim of disinfestation/disinfection, provided that any residue, if present, does not influence the assay. It is the responsibility of the user to check for such antagonism and/or inhibition by analysis, sample spiking, or experimental comparisons.

It is not validated for seed treated with protective chemicals or biological substances. If treated seed is tested, the user is responsible for determining empirically (through analysis, sample spiking, or experimental comparisons) whether the protective chemicals or biological substances have an effect on the method results.

METHOD EXECUTION

When pre-screening a seed lot using the ELISA test, it is strongly recommended that the best practices developed by ISHI-Veg for *ELISA Assays in Seed Health Tests* (see <https://www.worldseed.org/our-work/phytosanitary-matters/seed-health/ishi-veg-method-development/>) be followed to ensure process standardization and obtain valid results.

TMV, ToMV, ToMMV and ToBRFV are closely related and it has been shown that polyclonal antisera raised against one tobamovirus may show cross-reactivity with one or more of the other tobamoviruses. Commercial antisera specifically raised against ToMMV and ToBRFV may not yet be available; the antisera for TMV and ToMV may be used provided their reactivity with ToMMV and ToBRFV is verified. For PMMoV, more distantly related to the other four viruses, a specific antiserum must be used.

Note: The ELISA seed extract can also be used to perform the SE-qPCR in the ISHI-Veg test: *Detection of Infectious ToBRFV in Tomato and Pepper Seed*.

SAMPLE AND SUB-SAMPLE SIZE

The recommended minimum sample size is 3,000 seeds with a maximum sub-sample size of 250 seeds.

Protocol for detecting infectious tobamoviruses in pepper seed

ENZYME-LINKED IMMUNOSORBENT ASSAY (ELISA)

1. General Requirements

If a different seed extraction buffer is used, it must be verified in a comparison using uniform positive control material that it does not lead to a reduction in the number of lesions obtained.

2. Extraction of the virus from the seed

2.1. Grind 250 seeds in each sub-sample in 20 ml of Phosphate Buffered Saline (PBS), the seed extraction buffer in Table 2. Process seed extracts within 4 hours after grinding.

3. ELISA

3.1. Run a double-antibody-sandwich (DAS-)ELISA on the extracts (Albrechtsen, 2006; Clark and Adams, 1977).

LOCAL LESION ASSAY

Material

Assay plants (<i>Nicotiana glutinosa</i> or <i>Nicotiana tabacum</i> cv. Xanthi NN)	Carborundum (320 mesh grit powder, Fisher Scientific or equivalent)
Controls	Seed extraction buffer
Gloves or alkaline soap and water	Protective mask

Controls

Table 1: Description of controls

Positive Process Control (PPC)	A well-characterized positive seed lot
	Pea seed flour and powdered TMV, ToMV, ToMMV or ToBRFV infected leaves from a systemic host ground together in a seed extraction buffer (5 mL per 0.5 gr). The mix should be ground into a fine flour for efficient virus extraction
	Liquid extract of TMV, ToMV, ToMMV or ToBRFV infected leaves of Solanaceous hosts sufficiently diluted in PBS seed extraction buffer
Negative Process Control (NPC)	Seed extraction buffer
	Pepper seed free of tobamovirus

Seed extraction buffer

Table 2: Phosphate Buffered Saline (PBS) - pH 7.2 – 7.4 per liter

Sodium chloride (NaCl)	8.0 g
Disodium hydrogen phosphate (Na ₂ HPO ₄)	1.15 g
Potassium dihydrogen phosphate (KH ₂ PO ₄)	0.2 g
Add de-ionized water up to 1 liter, adjust pH and autoclave buffer at 121 °C, 15 psi for 15 minutes	

1. General Requirements

- Seed extracts and controls must be prepared at the same time, under the same laboratory conditions and stored under the same conditions.
- Seed extracts and controls must be stored at 4°C until the assay begins. When performed after pre-screening ELISA it is strongly recommended to perform the local lesion assay within 24 hours after seed-extraction.
- The final results of the local lesion assay must be validated through comparison of the results given by both controls.

2. Extraction of the virus from the seed

- 2.1. Grind 250 seeds in each sub-sample in 20 ml of the seed extraction buffer PBS.
- 2.2. Process seed extracts within 4 hours after grinding.

3. Controls

- 3.1. Choose one of the PPCs and NPCs listed.
- 3.2. Depending on the amount of infectious virus in the PPC, adjust the quantity used to induce development of countable local lesions.

4. Inoculation and incubation of *Nicotiana* assay plants

- 4.1. Raise assay plants (*Nicotiana glutinosa* or *Nicotiana tabacum* cv. Xanthi NN) under sufficient light intensity at a temperature of 20 – 25°C.
 - Choose plants with 4-5 true leaves (6-7 weeks after sowing) with high turgor. Do not use flowering or old plants (Takahashi, 1975); see Figure 2.
 - The primary leaf (oldest true leaf) should not be used as it is different in shape / texture / thickness and therefore less susceptible.
 - Leaves of tobacco plants affected by downy mildew are significantly less sensitive. Inoculated plant must have no visible symptoms and be free of disease (Koenraad H. personal communication).
- 4.2. Dust the leaves with the appropriate quantity of carborundum (320 mesh grit powder, Fisher Scientific or equivalent) so that there is a very fine layer on the leaf surface. Wear a protective mask.



Figure 2. Overview of *Nicotiana tabacum* cv. Xanthi NN plants in different growth stages (two pictures taken per plant; pot diameter 14 cm)

4.3. The entire surface of two (nearly) fully expanded consecutive leaves of two plants should be inoculated with each seed extract, a total of 4 leaves for each sub-sample.

4.4. Place a drop of inoculum (100-200 μ l) onto the leaf. Smear the drop with gloved fingers by applying constant, slight pressure but avoid damaging the leaf.

Note: Work with gloves and change them between samples or clean hands thoroughly between samples by using alkaline soap or equivalent and then rinse with water to remove residues.

4.5. Rinse the plants with tap water a few minutes after inoculation.

4.6. Inoculate plants with the PPC and NPC in a similar manner to the ones used for the seed extracts.

4.7. Incubate the plants for 5-7 days under controlled conditions at 20 – 25 °C and provide at least 12 hours of light.

Note: The hypersensitive reaction of tobacco plants, the development of local lesions, may not be expressed at temperatures higher than 28°C (Samuel, 1931; Takahashi, 1975; Weststeijn, 1981 and Kiraly et al., 2008) and may vary for different tobamoviruses.

5. Evaluation of inoculated plants

5.1. Evaluate the inoculated leaves for local lesions (see Figure 3).




Figure 3. Local lesions on tobacco indicator plants
(a) *Nicotiana glutinosa*, (b) *Nicotiana tabacum* cv. Xanthi NN

5.2. If there are many local lesions, a comparison with the PPC and NPC will readily reveal if most, or all, the lesions observed are authentic virus lesions or might be caused by, for instance, rubbing (rubbing damage).

5.3. Optional: If the number of lesions is low, confirm they are a true virus infection and not an artefact (leaf damage during mechanical inoculation, use of pesticides, etc.) by cutting out the suspect lesion, crushing it in a small amount of the seed extraction buffer and inoculating two leaves of two assay plants again. Lesions caused by virus infection contain sufficient amounts of infectious virus to produce multiple lesions in this confirmation test (Hadas, 2000).

REFERENCES

- Albrechtsen, S.E. (2006). Testing methods for seed-transmitted viruses: principles and protocols. Wallingford, UK: CABI Publishing.
- Clark, M.F. and Adams, A.N. (1977). Characteristics of the microplate method of enzyme-linked immunosorbent assay for the detection of plant viruses. *Journal of General Virology* 34: 475-483.
- Hadas, R. (2000). A report of comparative test for Tobamoviruses in pepper seeds. ISHI-Veg Research Report 1-2000-pepper-tobamo. Nyon, Switzerland: International Seed Federation.
- Kiraly, L., Hafez, Y. M., Fodor, J. and Kiraly, Z. (2008). Suppression of tobacco mosaic virus-induced hypersensitive-type necrotization in tobacco at high temperature is associated with down regulation of NADPH oxidase and superoxide and stimulation of dehydroascorbate reductase. *Journal of General Virology* 89, 799-808.
- Li, R., Gao, S., Fei, Z. and Ling, K.S. (2013). Complete genome sequence of a new tobamovirus naturally infecting tomatoes in Mexico. *Genome announcements*, 1(5): e00794-13.
- Luria, N., Smith, E., Reingold, V., Bekelman, I., Lapidot, M., Levin, I., Elad, N., Tam, Y., Sela, N., Abu-Ras, A. and Ezra, N. (2017). A new Israeli Tobamovirus isolate infects tomato plants harboring Tm-2² resistance genes. *PLoS one*, 12 (1), p.e0170429.
- Salem, N., Mansour, A., Ciuffo, M., Falk, B.W. and Turina, M. (2016). A new tobamovirus infecting tomato crops in Jordan. *Archives of virology*, 161 (2), pp. 503-506.
- Samuel, G. (1931). Some experiments on inoculating methods with plant viruses and on local lesions. *Annals of Applied Biology* 18, 494-507.
- Sui, X., Zheng, Y., Li, R., Padmanabhan, C., Tian, T., Groth-Helms, D., Keinath, A.P., Fei, Z., Wu, Z. and Ling, K.S. (2017). Molecular and Biological Characterization of Tomato mottle mosaic virus and Development of RT-PCR Detection. *Plant disease*, 101 (5), pp. 704-711.
- Takahashi, T. (1975). Studies on viral pathogenesis in plant hosts VIII. Systemic virus invasion and localization of infection on "Samsun NN" tobacco plants resulting from tobacco virus infection. *Phytopathology*, Z. 84, 75-87.
- Weststeijn, E. A. (1981). Lesion growth and virus localization in leaves of *Nicotiana tabacum* cv. Xanthi nc. after inoculation with tobacco mosaic virus and incubation alternately at 22°C and 32°C. *Physiological Plant Pathology* 18, 357-368.